

NATURAL AND ARTIFICIAL NUCLIDES IN SALESÓPOLIS RESERVOIR BY GAMMA SPECTROMETRY

Paulo S. C. Silva, Renato Semmler, Guilherme S. Zahn, Flávio R. Rocha, Sandra R. Damatto, Déborah I. T. Fávoro

Instituto de Pesquisas Energéticas e Nucleares (IPEN / CNEN - SP)
Av. Prof. Lineu Prestes, 2242, Cidade Universitária
05508-000 São Paulo, SP
pscilva@ipen.br
rsemmler@ipen.br
gzahn@ipen.br
flavio@baquara.com
damatto@ipen.br
defavaro@ipen.br

ABSTRACT

Natural radioactivity is ubiquitous in the environment mainly due to the presence of the nuclides from the uranium and thorium series and ^{40}K . Although in the South Hemisphere nuclear tests have been fewer in number than that in the North, artificial radionuclides can also be found spread at ground level. In this study, the activity concentrations of natural nuclides from the uranium and thorium series, ^{40}K and the artificial ^{137}Cs were determined in a sediment core with 42 cm depth collected in the middle of the Salesópolis reservoir, located in the Metropolitan Region of São Paulo city (SPMR). The Usina Parque Rio Tietê reservoir belongs to the Alto do Tietê system for the capture, storage and treatment of water for SPMR. Therefore, the quality of the water and sediments of this dam is of great importance. The activity concentrations were measured by gamma spectrometry. Samples were measured and saved at regular intervals at a maximum of 160 000 seconds. The gross area were determined for each peak and plotted against time and the counting rate was obtained by the slope of the curve. Background and reference materials were also counted and treated in the same way. Results showed that ^{226}Ra varied from 45 to 116 Bq kg^{-1} ; ^{228}Ra , from 80 to 165 Bq kg^{-1} ; ^{40}K , from 155 to 1 187 Bq kg^{-1} and ^{137}Cs varied from 0.3 to 7 Bq kg^{-1} . The methodology applied for determining low levels of ^{137}Cs in sediment proved to be efficient and reproducible.

1. INTRODUCTION

Environmental radioactivity occurs mainly due to the presence of natural radionuclides of the uranium and thorium series and ^{40}K in soil, rocks, water and air along with artificial nuclides also spread in the biosphere after nuclear accidents and nuclear weapon tests [1]. Although in the South Hemisphere nuclear tests have been fewer in number than that in the North, artificial radionuclides can also be found spread at ground level [2, 3]. The distribution of the radionuclides once released in the environment to the different compartments depends on their different geochemical behaviors, as they are subjected to a number of natural processes such as dissolution, adsorption, complexation, and oxidation/reduction.

The natural and man made radionuclides may eventually be accumulated in sediments due to processes like weathering and terrestrial minerals and rocks recycling, rainfall and other depositional phenomena such as gravitational settling and precipitation [4]. Sediments, in turn, can act as sink for these elements or release them to the surrounding water depending on

the physico-chemical conditions. This paper aims to evaluate the distribution of the radionuclides belonging to the U and Th series, ^{40}K and the artificial ^{137}Cs in a sediment core of the Salesópolis reservoir by gamma spectrometry.

Salesópolis is located in the São Paulo city Metropolitan Region (SPMR). The Usina Parque Rio Tietê (Salesópolis) reservoir belongs to the Alto do Tietê system for the capture, storage and treatment of water for the SPMR. Therefore, the quality of the water, as well as of sediments of this dam is of great importance.

2. MATERIAL AND METHODS

A sediment core named T1B with 42 cm depth was collected in the Salesópolis dam in 2014 with a manual core sampler under a water column of approximately one and a half meters. In the laboratory, the core was sliced each 3 cm totalizing 14 samples. Each sample was then dried in a ventilated oven at a temperature of 45 °C and passed in a sieve of 2 mm aperture. After that, approximately 25 g of samples were weighted in plastic containers of 5 cm diameter and 1.5 cm of height, sealed and let to wait at least 30 days for the radioactive equilibrium between ^{226}Ra and its daughters to be reached.

Gamma activity was measured in a EG&G Ortec spectrometer with relative efficiency of 60%, and resolution of 1.16 and 2.25 keV for the 122 and 1332 keV photopeaks of ^{57}Co and ^{60}Co , respectively. In order to diminish the background (BG) contribution the detector is shielded with a sequential layer of lead and copper combined protection.

The activity concentration of ^{226}Ra was determined by the mean activity measurement of its daughter's photopeaks in 351 keV (^{214}Pb), 609 and 1120 keV (^{214}Bi). The activity concentration of ^{228}Ra was determined by the gamma transitions in 338 keV, 911 keV (^{228}Ac), 238 keV (^{212}Pb) and 727 keV (^{212}Bi). Potassium-40 and ^{137}Cs were determined directly by means of their gamma transition photopeaks in 1460 keV and 661 keV, respectively.

The efficiency curve determination was performed by using the reference materials IAEA-RGU (uranium ore diluted with floated silica powder), IAEA-RGTh (thorium ore diluted with floated silica powder), IAEA-K (high purity potassium sulphate) and IAEA-375 (radionuclides and trace elements in soil) in the same geometry as the samples for the same gamma transitions photopeaks to be determined. All these reference material have activity sufficiently high to give a statistical error of less than 1% in the considered counting time. No self-attenuation correction was applied since the energy of all determined nuclides was higher than 200 keV.

For the data acquisition, the processing software was programmed to save the counting files from 100 000 s to 160 000 s in intervals of 10 000 s. A background measurement was also done intercalating each two samples counting. This BG counts was subtracted from the samples counting [5].

For each sample and respective BG a plot of gross counts per time was generated to obtain the curve slope. Figures 1 and 2 exemplify the procedure for ^{137}Cs determination and that was then repeated for each radionuclide to be determined.




Figure 1: Plot of the background counts per time (in seconds) in the 661 keV region of interest




Figure 2: Plot of the counts per time (in seconds) in the 661 keV region of interest for a given sample

Calculation of the activity concentration was done according to the formula presented in equation 1.

$$A_i = \frac{C_i - C_{bgi}}{m_a \times \varepsilon_i} \quad (1)$$

where:

- A_i is the activity concentration related to the photopeak i (Bq kg^{-1})
- C_i is the slope of the curve obtained for the photopeak i in the sample
- C_{bgi} is the slope of the curve obtained for the photopeak i in the background
- m_a is the sample mass (kg)
- ε_i is the counting efficiency for the photopeak of interest.

The uncertainties were determined by error propagation, with the uncertainties of C_i and C_{bgi} being that obtained from the slope of the adjusted curve for each photopeak considered.

3. RESULTS AND DISCUSSION

Table 1 shows the activity concentration measured in the reference material IAEA-414, IAEA-SD-N-2 and IAEA-375 and it is possible to see the good agreement between the measured values and the certified ones.

Table1: Results obtained for the reference materials IAEA-327, IAEA-414 and IAEA-SD-N-2, in Bq kg⁻¹ (certified values in parenthesis)

	Reference material	Obtained value	Certified value
²²⁶ Ra	IAEA-327	43 ± 8	38.7 ± 5.2
²²⁸ Ra	IAEA-327	40 ± 7	38.7 ± 4.7
⁴⁰ K	IAEA-327	641 ± 41	621 ± 52
	IAEA-414	406 ± 26	481 ± 16.5
	IAEA-SD-N-2	218 ± 14	220 ± 16
¹³⁷ Cs	IAEA-327	28 ± 7	24.9 ± 1.8
	IAEA-414	5 ± 1	5.18 ± 0.1
	IAEA-SD-N-2	0.9 ± 0.2	0.8 ± 0.5

3.1 Natural radionuclides

Table 2 presents the values of the determined radionuclides in the samples obtained in the analyzed core. It is possible to see that radium isotopes are in higher concentrations than that reported for the global soil average [6] of 32 Bq kg⁻¹ for ²²⁶Ra and 45 Bq kg⁻¹ for ²²⁸Ra (considered in equilibrium with ²³²Th). For ⁴⁰K, only the first 12 cm are below the global average of 420 Bq kg⁻¹ [6] and the activity concentrations tend to rise with the depth.

Figure 3 shows that the radium isotopes have almost the same behavior with depth, with the lowest values observed between 9 and 12 cm depth. Higher activity concentrations were also observed for both nuclides in the base of the core. Nevertheless, ²²⁸Ra tend to be enriched over ²²⁶Ra in the top layers with the enrichment being less pronounced towards the core base as depicted by the ²²⁸Ra/²²⁶Ra ratio showed in Figure 4 probably indicating a change in the sediment source and mineralogy.

Few data are available on the activity concentrations of of natural radionuclides in Brazilian reservoirs. Concentrations lower than those observed in this study, around 20 Bq kg⁻¹, were determined for ²²⁶Ra in the Apipucos Reservoir, Recife, Brazil [7]. In sediments of Cubatão river, ²²⁶Ra activity concentration varied in the range of 28 to 80 Bq kg⁻¹ [8] and the activity concentrations of ²²⁶Ra and ²²⁸Ra were in the range of 26 to 37 Bq kg⁻¹ and 59 to 74 Bq kg⁻¹, respectively [9]. Considering aquatic environments with saline water, levels of ²³⁸U (²²⁶Ra), ²³²Th and ⁴⁰K of 21, 28 and 426 Bq kg⁻¹ were determined [10] in the shelf and upper slope of

Southeast Brazil. Also levels of ^{226}Ra in the range of 4.2 to 28.5 were determined in the Cananéia- Iguape system [11]. Mean activity concentrations of 24 Bq kg⁻¹ for ^{226}Ra , 44 Bq kg⁻¹ for ^{232}Th and 678 Bq kg⁻¹ for ^{40}K were found in the Piraquara de Fora Inlet located in Ribeira Bay within the Ilha Grande Bay in Angra dos Reis, Rio de Janeiro State [3]. It can be seen that the values obtained in this study for the Salesópolis reservoir are generally higher than those reported in literature and that sediments of salty waters generally presents lower activity concentrations.

Table 2: Activity concentrations, in Bq kg⁻¹, of measured radionuclides in the sediment core samples

Depth (cm)	^{226}Ra	^{228}Ra	^{40}K	^{137}Cs
0-3	74±13	126±31	300±19	5±1
3-6	67±12	132±33	155±10	2±1
6-9	72±13	105±26	272±17	7±3
9-12	45±8	80±20	179±11	0.3±0.1
12-15	67±12	113±20	680±43	3±1
15-18	79±14	133±33	807±52	4±2
18-21	65±11	108±27	856±55	3±1
21-24	72±13	116±29	843±54	3±1
24-27	70±12	103±26	830±53	4±2
27-30	79±11	122±30	978±63	4±2
30-33	83±14	130±32	1 026±66	4±2
33-36	75±13	122±30	940±60	4±2
36-39	89±15	124±31	950±61	5±2
39-42	116±20	165±41	1 187±76	4±2




Figure 3: Distribution of ^{226}Ra and ^{228}Ra with the depth in the analyzed core




Figure 4: Distribution of $^{228}\text{Ra}/^{226}\text{Ra}$ isotopic ratio with depth

Table 3 shows a compilation of activity concentrations for radium isotopes, ^{40}K and ^{137}Cs , found in sediments of reservoirs located in North Hemisphere. For the natural radionuclides it can be seen that the levels are closer to those observed for the global average [6] and present a mean narrower those observed for the Brazilian reservoir and the results obtained in this work.

Higher activity concentrations for natural radionuclides may be related to geological and geochemical conditions such as the rock and soil basement and leaching by an intense rainfall system.

3.2 Artificial radionuclide ^{137}Cs

The distribution of ^{137}Cs with depth is shown in Figure 5. The lowest activity concentration was observed at 9 – 12 cm depth and the highest was determined in the range of 6 to 9 cm, although it is unlikely that this peak corresponds to the maximum fallout of Cs. The observed variation in the top layers may be related to the grain size and organic matter distribution [18].

Table 3: Activity concentrations, in Bk kg^{-1} , of natural and artificial radionuclides in sediment samples of reservoirs from selected locations in the North Hemisphere

	^{226}Ra	^{228}Ra	^{40}K	^{137}Cs	Reference
Spain	58 - 94	59 - 116	535 - 823	1.7 - 60.2	[12]
India	4 - 26	5.5 - 19.9	250 - 558	3.6 - 54.5	[13]
Tunisia	23 - 29.8	40 - 52		0.9 - 3.9	[14]
Serbia	8 - 29	2.7 - 45		0.8 - 140	[15]
China	31.2 - 37.5	56.5 - 79.8	179.4 - 214.9		[16]
Poland				89 - 865	[17]
Poland				30 - 243	[18]
Turkey				1 - 101	[19]




Figure 5: Distribution of ^{137}Cs with the depth in the analyzed core

The most probable peak of ^{137}Cs activity concentration corresponding to the fallout of 1963-1964 can be observed in the depth of 36 – 39 cm. As for the natural radionuclides, few activity concentrations were reported for ^{137}Cs in reservoir sediments. Considering Brazilian sediment of other aquatic environments, levels of this radionuclide in the range of 0.2 to 6.1 Kq kg^{-1} were found in the Cananéia-Iguape system [11]; activity concentration of $0.5 \pm 0.2 \text{ Bq kg}^{-1}$ for ^{137}Cs was determined in the Piraquara de Fora Inlet [3]; values around 3 Bq kg^{-1} were found in the sediments of Guajar Bay, in Par State [2] and a maximum of 4.7 Bq kg^{-1} was reported [20] for the Brazilian coast. The values obtained in this study are in good agreement with those observed in the references cited above, and all the highest values found in Brazilian sediments were much lower than those found in the North Hemisphere, as shown in Table 3. Closer values to the ones reported here were found in a South African reservoir with activity concentrations around 10 Bq kg^{-1} [21].

4. CONCLUSIONS

The methodology applied for determining low levels of ^{137}Cs in sediment proved to be efficient and reproducible. The levels of natural radionuclides found in the sediment core of Salesópolis dam are higher than the world average. These activities can be related to the local geochemistry associated with intense rainfall. Results showed a tendency of raising values in the core base possibly indicating a variation in the mineralogical features as shown by the decrease in the $^{228}\text{Ra}/^{226}\text{Ra}$ ratio.

The activity concentrations of ^{137}Cs present a great variation in the upper layers probably related to the granulometry distribution and organic matter content. The values found are in accordance with the ones observed for other Brazilian aquatic environments. Nevertheless, more studies are necessary to better understand the observed variation as well as the role of local conditions in the observed values.

ACKNOWLEDGMENTS

Authors are thankful to the Fundação de Amparo à Pesquisa do Estado de São Paulo – FAPESP – for the financial support (FAPESP Process n^o: 2014/20805-6).

REFERENCES

1. IAEA. *Radiation, People and Environment*. International Atomic Energy Agency, Viena, Austria, (2004).
2. P. A. L. Ferreira, P. A. Neves, R. C. L. Figueira, Evaluation of recent sedimentation rates in Guajará Bay (in Brasilina coast) with ^{210}Pb and ^{137}Cs . *International Nuclear Atlantic Conference - INAC 2013*, Recife, PE, Brazil, November 24 - 29, (2013).
3. F. M. Carvalho, D. C. Lauria, F. C. A. Ribeiro, R. T. Fonseca, S. S. Peres, N. S. F. Martins, “Natural and man-made radionuclides in sediments of an inlet in Rio de Janeiro State, Brazil”, *Marine Pollution Bulletin*, **107**, pp. 269 - 276 (2016).
4. F. Caridi, S. Marguccio, A. Belvedere, G. Belmusto, “Measurements of Gamma Radioactivity in River Sediment Samples of the Mediterranean Central Basin”, *American Journal of Condensed Matter Physics*, **5(3)**, pp. 61- 68 (2015).
5. R. C. L. Figueira, R. L. N. Silva, A. M. G. Figueiredo, I. I. L. Cunha, Instrumental analysis by gamma spectrometry of low level caesium-137 in marine samples. *Proceedings of an International Conference*, Goiania, Brazil, 26 - 31 october, pp. 329 – 337 (1997).
6. UNSCEAR. *Sources and Effects of Ionizing Radiation*. Volume I: Effects. UNSCEAR 2008 Report. United Nations Scientific Committee on the Effects of Atomic Radiation, Report to the General Assembly, with scientific annexes. United Nations sales publication E.00.IX.4. United Nations, New York (2010).
7. V. L. B. Souza, K. R. G. Rodrigues, E. H. Pedroza, R. T. Melo, V. L. Lima, C. A. Hazin, M. G. O. Almeida, R. K. Nascimento, “Sedimentation Rate and ^{210}Pb Sediment Dating at Apipucos Reservoir, Recife, Brazil”, *Sustainability*, **4**, pp. 2419 - 2429 (2012).

8. L. M. Sanders, C. J. Sanders, W. Luiz-Silva, W. Machado, E. V. Silva-Filho, S. R. Patchineelam, “Anthropogenic source assessment of ^{226}Ra and ^{210}Pb in a sediment core from the Cubatão River estuary (SE Brazil)”, *Journal of Radioanalytical and Nuclear Chemistry*, **287(3)**, pp. 729 - 732 (2011).
9. P. S. C. Silva, B. P. Mazzilli, D. I. T. Fávoro, “Distribution of radionuclides and elements in Cubatão River sediments”, *Journal of Radioanalytical and Nuclear Chemistry*, **269 (3)**, pp. 767 - 771 (2006).
10. L. M. Cordero, R. C.L. Figueira, M. M. Mahiques, M. G Tessler, “Natural radionuclides distribution in the shelf and upper slope off Southeast Brazil”. *International Nuclear Atlantic Conference – INAC*, Rio de Janeiro, RJ, Brazil, September 27 to October 2, pp. 1 – 7 (2009).
11. R. T. Saito, *Radionuclídeos (^{210}Pb , ^{226}Ra , ^{210}Po e ^{137}Cs) no sistema costeiro Canaéia-Iguape: estudos ambientais*. Doctorate Thesis IPEN, (2002). (in portuguese).
12. A. Baeza, J. Guillén, M. A. Ontalba Salamanca, A. Rodríguez, F. J. Ager, “Radiological and multi-element analysis of sediments from the Proserpina reservoir (Spain) dating from Roman times”, *Journal of Environmental Radioactivity*, **100**, pp. 866 - 874 (2009).
13. S. Sugandhi, V. M. Joshi, P. M. Ravi, “Studies on natural and anthropogenic radionuclides in sediment and biota of Mumbai Harbour Bay”, *J Radioanal Nucl Chem*, **300**, pp. 67 - 70 (2014).
14. A. B. Slimane, D. Raclot, O. Evrard, M. Sanaa, I. Lefèvre, M. Ahmadi, M. Tounsi, C. Rumpel, A. B. Mammou, Y. Bissonais, “Fingerprinting sediment sources in the outlet reservoir of a hilly cultivated catchment in Tunisia”, *J Soils Sediments*, **13**, pp. 801 - 815 (2013).
15. R. Tošić, D. J. Todorović, S. S. Dragičević, I. S. Bikit, S. Forkapić, B. Blagojević, “Radioactivity and measurements of sediment deposition rate of the Drenova Reservoir (B&H)”, *Nuclear Technology & Radiation Protection*, **27 (1)**, pp. 52 - 56 (2012).
16. X. Lu, H. Pan, C. Ren, L. Yang, “Natural radioactivity in reservoir sediment near an industrial park of northwest China”, *J Radiol Prot*, **36**, pp. N26 - N33 (2016).
17. K. Szarlowicz, W. Reczynski, R. Misiak, B. Kubica, “Radionuclides and heavy metal concentrations as complementary tools for studying the impact of industrialization on the environment”, *J Radioanal Nucl Chem*, **298**, pp. 1323 - 1333 (2013).
18. K. Szarlowicz, B. Kubica, “ ^{137}Cs and ^{210}Pb radionuclides in open and closed water ecosystems”, *J Radioanal Nucl Chem*, **299**, pp. 1321 - 1328 (2014).
19. F. Kuilahcıa, Z. Şen, “Multivariate statistical analyses of artificial radionuclides and heavy metals contaminations in deep mud of Keban Dam Lake, Turkey”, *Applied Radiation and Isotopes*, **66**, 236 - 246 (2008).
20. I. R. Santos, W. C. Burnett, J. M. Godoy, “Radionuclides as tracer of coastal processes in Brazil: review, synthesis, and perspectives”, *Brazilian Journal of Oceanography*, **56(2)**, 115 - 131, (2008).
21. I. D.L. Foster, J. Boardman, J. Keay-Bright, “Sediment tracing and environmental history for two small catchments, Karoo Uplands, South Africa”, *Geomorphology*, **90**, 126 - 143 (2007).