

**RECENT SEDIMENTATION RATES AND TRACE ELEMENTS DETERMINED IN CORES  
FROM PANTANAL, MATO GROSSO DO SUL, BRASIL**

**Sandra Regina DAMATTO<sup>1</sup>, Barbara Paci MAZZILLI<sup>1</sup>, Déborah Inês Teixeira FÁVARO<sup>2</sup>,  
Arnaldo SAKAMOTO<sup>3</sup>**

1) Laboratório de Radiometria Ambiental – IPEN, São Paulo, Brasil

2) Laboratório de Análise por Ativação – IPEN

3) Universidade Federal do Mato Grosso do Sul

E-mail: [damatto@ipen.br](mailto:damatto@ipen.br)

Pantanal, in the southwest part of Brazil, is one of the world's largest freshwater wetland. This natural ecosystem has been affected due to urban contamination, irregular use of the land, tourism without control, excessive agricultural defensive utilization, etc.

In order to verify possible changes in this environment, a study was established in Pantanal da Nhecolândia, Mato Grosso do Sul, Brazil.

In 2006 three sediment cores were collected in two small ponds in Nhumirim farm, Salina do Meio, Salina Pedra do Sol and Salina da Ponta. The sedimentation rate was determined using the <sup>210</sup>Pb dating method.

The elements As, Ba, Br, Ce, Co, Cr, Cs, Eu, Fe, Hf, La, Lu, Na, Nd, Rb, Sb, Sc, Sm, Ta, Tb, Th, U, Yb, Zn and Zr were determined by instrumental neutron activation analysis technique – NAA.

Their concentrations ranged from mg kg<sup>-1</sup> to %. For validation methodology the reference materials Buffalo River Sediment (NIST SRM 2704) and Soil-7 (IAEA) were analyzed.

The majority of the elements and rare earths analyzed showed concentrations lower when compared with shale and earth crust values. Factorial analysis, mode R, Cluster analysis and principal component analysis were applied for the data interpretation.