

# Decoupled cooling and exhumation signals in subduction orogens: An example from the Argentinean Pampean Ranges

Mauricio Parra<sup>1</sup>, Miguel Ezpeleta<sup>2</sup>, Gilda Collo<sup>2</sup>, Cecilia Wunderlin<sup>2</sup>, Edward Sobel<sup>3</sup>,  
Ángeles Borrego<sup>4</sup>, and Frederico Genezini<sup>5</sup>

<sup>1</sup>Instituto de Geociências, Universidade de São Paulo, Brazil

<sup>2</sup>Centro de Investigaciones en Ciencias de la Tierra (CICTERRA), Universidad Nacional de Córdoba, Argentina

<sup>3</sup>Institut fuer Geowissenschaften, Universitaet Potsdam

<sup>4</sup>Instituto de Ciencia y Tecnología del Carbono (INCAR-CSIC), Oviedo, Spain

<sup>5</sup>Instituto de Pesquisas Energéticas e Nucleares

November 26, 2022

## Abstract


Reconstructing thermal histories in thrust belts using thermochronometry is a widely used method to infer the age and rates of thrusting, a precondition to understanding the driving mechanisms of orogenesis. Along a thrust sheet, the time and temperature conditions at the switch between heating and cooling retrieved from thermal modeling are commonly interpreted as the onset of thrust-induced exhumation associated with thrustbelt development. In subduction orogens such as the Andes, this interpretation can be challenged by the intrinsic changes in basal heat flow imposed by changes in subduction regimes. We document a case in the northwestern Sierras Pampeanas in the Argentinean Central Andes in which independent constraints on the onset late Cenozoic thrusting derived from structural cross-cutting relationships allow us to explore alternative causes for Cenozoic cooling signals. Located at  $\sim 31^\circ\text{S}$  Lat, the Villa Unión-Ischigualasto basin hosts a composite stratigraphic record associated with Triassic rifting developed onto the Paleozoic substratum of western Gondwana and the overlying Meso-Cenozoic foreland basin record. A multi-method approach including apatite fission-track, apatite and zircon (U-Th)/He analyses, vitrinite reflectance and clay mineralogy carried out along three stratigraphic profiles, and inverse thermal modeling reveals the thermal history patterns and allows inferring its triggering mechanisms. Despite an up to 5 km-thick Cenozoic overburden and unlike previously thought, the thermal peak in the basin is not due to Cenozoic burial but occurred in the Triassic, associated with an abnormally high heat flow of up to  $90 \text{ mWm}^{-2}$  and less than 2 km of burial, which heated the base of the Triassic strata to  $\sim 160^\circ\text{C}$ . Following exhumation, attested by the development of an unconformity between the Triassic and Late-Cretaceous-Cenozoic sequences, Cenozoic re-burial increased the temperature to  $\sim 110^\circ\text{C}$  at the base of the Triassic section and only  $\sim 50^\circ\text{C}$  4 km upsection, suggesting a dramatic decrease in the thermal gradient. The onset of Cenozoic cooling from those conditions occurred between  $\sim 10$  and 8 Ma, approximately 5 My before the onset of thrusting that has been independently documented by exceptionally well preserved stratigraphic-cross-cutting relationships. We argue that the onset of cooling is associated with lithospheric refrigeration following a decrease in the angle of subduction of the Nazca slab, leading to the eastward displacement of the asthenospheric wedge beneath the South American plate. Our study places time and temperature constraints on an idea that has been previously discussed in the region and calls for a careful interpretation of exhumation signals in thrustbelts inferred from thermochronology only.




Mauricio Parra<sup>1\*</sup>, Miguel Ezpeleta<sup>2</sup>, Gilda Collo<sup>2</sup>, Cecilia A. Wunderlin<sup>2</sup>, Edward Sobel<sup>3</sup>, Ángeles G. Borrego<sup>4</sup> and Federico Genezini<sup>5</sup>

(1) Instituto de Geociências, Universidade de São Paulo, Brazil, (2) Centro de Investigaciones en Ciencias de la Tierra (CICTERRA), Universidad Nacional de Córdoba, Argentina, (3) Institut für Geowissenschaften, Universität Potsdam, Germany, (4) Instituto de Ciencia y Tecnología del Carbono (INCAR-CSIC), Oviedo, Spain, (5) Centro do Reator Nuclear de Pesquisa, Instituto de Pesquisas Energéticas e Nucleares (IPEN), Brazil


## INTRODUCTION




## 1. STRATIGRAPHY AND MULTI-METHOD PALEOTHERMOMETRY




## 2. THERMOCHRON AGES AND THERMAL MODELS




## 3. MULTI-SAMPLE THERMAL MODELING



## 4. FORWARD THERMAL MODELING



## 5. COOLING VS EXHUMATION SIGNALS



## Example of cooling and exhumation signal decoupled in subduction orogens, call for a rigorous interpretation of cooling paths in Andean fold-and-thrust belts.

## ACKNOWLEDGEMENTS

Funding by the Sao Paulo Research Foundation (FAPESP, Brazil) and Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET, Argentina) binational project 2016/50441-1 "Multi-method Paleothermometry in Sedimentary Basins".

REFERENCES  
Fosdick et al. (2015), Faulting and erosion in the Argentine Precordillera during changes in subduction regime: Reconciling bedrock cooling and detrital records, *Earth and Planetary Science Letters* 432, 73-83.  
Jordan et al. (2001), Unsteady and spatially variable evolution of the Neogene Bermejo foreland basin, Argentina, *Journal of South American Earth Sciences* 14, 775-798.  
Ortiz et al. (2015), Active deformation in the northern Sierra de Valle Fértil, Sierras Pampeanas, Argentina, *Journal of South American Earth Sciences* 64, 339 - 350.